

BETA-EXPANSIONS OF RATIONAL NUMBERS IN QUADRATIC PISOT BASES

TOMÁŠ HEJDA AND WOLFGANG STEINER

ABSTRACT. We study rational numbers with purely periodic Rényi β -expansions. For bases β satisfying $\beta^2 = a\beta + b$ with b dividing a , we give a necessary and sufficient condition for $\gamma(\beta) = 1$, i.e., that all rational numbers $p/q \in [0, 1)$ with $\gcd(q, b) = 1$ have a purely periodic β -expansion. A simple algorithm for determining the value of $\gamma(\beta)$ for all quadratic Pisot numbers β is described.

1. INTRODUCTION AND MAIN RESULTS

Rényi β -expansions [Rén57] provide a very natural generalization of standard positional numeration systems such as the decimal system. Let $\beta > 1$ denote the base. Expansions of numbers $x \in [0, 1)$ are defined in terms of the β -transformation

$$T: [0, 1) \rightarrow [0, 1), \quad x \mapsto \beta x - \lfloor \beta x \rfloor.$$

The expansion of x is the infinite string $x_1x_2x_3\cdots$ where $x_j := \lfloor \beta T^{j-1}x \rfloor$. For $\beta \in \mathbb{N}$, we recover the standard expansions in base β and the β -expansion of $x \in [0, 1)$ is eventually periodic (i.e., there exist p, n such that $x_{k+p} = x_k$ for all $k \geq n$) if and only if $x \in \mathbb{Q}$. This result was generalized to all Pisot bases by Schmidt [Sch80], who proved that for a Pisot number β the expansion of $x \in [0, 1)$ is eventually periodic if and only if x is an element of the number field $\mathbb{Q}(\beta)$. Moreover, he showed that when β satisfies $\beta^2 = a\beta + 1$, then each $x \in [0, 1) \cap \mathbb{Q}$ has a purely periodic β -expansion.

Akiyama [Aki98] showed that if β is a Pisot unit satisfying a certain finiteness property then there exists $c > 0$ such that all rational numbers $x \in \mathbb{Q} \cap [0, c)$ have a purely periodic expansion. If β is not a unit, then a rational number $p/q \in [0, 1)$ can have a purely periodic expansion only if q is co-prime to the norm $N(\beta)$. Many Pisot non-units satisfy that there exists $c > 0$ such that all rational numbers $\frac{p}{q} \in [0, c)$ with q co-prime to b have a purely periodic expansion. This stimulates for the following definition:

Definition 1.1. Let β be a Pisot number, and let $N(\beta)$ denote the norm of β . Then we define $\gamma(\beta) \in [0, 1]$ as the maximal c such that all $\frac{p}{q} \in \mathbb{Q} \cap [0, c)$ with $\gcd(q, N(\beta)) = 1$ have a purely periodic β -expansion. In other words,

$$\gamma(\beta) := \inf \left\{ \frac{p}{q} \in \mathbb{Q} \cap [0, 1) : \gcd(q, b) = 1, \frac{p}{q} \text{ has a not purely periodic expansion} \right\} \cup \{1\},$$

where $b = |N(\beta)|$ is the norm of β .

The question is how to determine the value of $\gamma(\beta)$. As well, knowing when $\gamma(\beta) = 0$ or 1 is of big interest. Values of $\gamma(\beta)$ for whole classes of numbers as well as for particular numbers have been given [Aki98, ABBS08, AS05, MS14, Sch80].

It is easy to observe that the expansion of x is purely periodic if and only if x is a periodic point of T , i.e., there exists $p \geq 1$ such that $T^p x = x$. The natural extension $(\mathcal{X}, \mathcal{T})$ of the dynamical system $([0, 1), T)$ (w.r.t. its unique absolutely continuous invariant measure) can be defined in an algebraic way, cf. §2.3. Several authors contributed to proving the following result: A point $x \in [0, 1)$ has a purely periodic β -expansion if and only if $x \in \mathbb{Q}(\beta)$ and its diagonal embedding lies in the natural extension domain $\mathcal{X}$. The quadratic unit case was solved by Hama and Imahashi [HI97], the confluent unit case by Ito and Sano [IS01, IS02]. Then Ito and Rao [IR05] resolved the unit case completely using an algebraic argument. For non-unit bases β , one has to consider finite

(p -adic) places of the field $\mathbb{Q}(\beta)$. This consideration allowed Berthé and Siegel [BS07] to expand the result to all (non-unit) Pisot numbers.

The first values of $\gamma(\beta)$ for two particular quadratic non-units were provided by Akiyama et al. [ABBS08]. Recently, Minervino and Steiner [MS14] described the boundary of $\mathcal{X}$ for quadratic non-unit Pisot bases. This allowed them to find the value of $\gamma(\beta)$ for an infinite class of quadratic numbers:

Theorem A ([MS14]). *Let β be the positive root of $\beta^2 = a\beta + b$ for $a \geq b \geq 1$ two co-prime integers. Then*

$$\gamma(\beta) = \begin{cases} 1 - \frac{(b-1)b\beta}{\beta^2 - b^2} & \text{if } a > b(b-1), \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $\gamma(\beta) = 1$ if and only if $b = 1$.

The purpose of this article is to generalize Theorem A to all quadratic Pisot numbers with norm $N(\beta) < 0$. (Note that when $N(\beta) > 0$, then β has a positive Galois conjugate $\beta' > 0$ and $\gamma(\beta) = 0$ by [Aki98, Proposition 5].) To this end, we define β -adic expansions (not to be confused with the Rényi β -expansions) similarly to p -adic expansions with $p \in \mathbb{Z}$, see also §2.4.

Definition 1.2. Let β be an algebraic integer. The β -adic expansion of $x \in \mathbb{Z}[\beta]$ is the unique infinite word $\mathbf{h}(x) := u_0 u_1 u_2 \dots$ such that $u_n \in \{0, 1, \dots, |N(\beta)| - 1\}$ and $x - \sum_{i=0}^{n-1} u_i \beta^i \in \beta^n \mathbb{Z}[\beta]$ for all $n \in \mathbb{N}$.

Theorem 1. *Let β be a quadratic Pisot number, root of $\beta^2 = a\beta + b$ with $a \geq b \geq 1$. Then*

$$\gamma(\beta) = \begin{cases} 0 & \text{if } \sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta') > \beta \text{ or } \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') < -1, \\ \beta - a & \text{if } \sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta') \in (2\beta - a - 1, \beta] \text{ and } \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \geq \beta - a - 1, \\ 1 + \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') & \text{otherwise,} \end{cases}$$

where $P_{u_0 u_1 u_2 \dots}(X) := \sum_{n \geq 0} u_n X^n$.

In many cases, we obtain the following direct formula (which we conjecture to be true for all $a \geq b \geq 1$):

Theorem 2. *Let β be a quadratic Pisot number, root of $\beta^2 = a\beta + b$ for $a \geq b \geq 1$. Suppose $a > \frac{1+\sqrt{5}}{2}b$ or $a = b$ or $a \perp b$. Then*

$$(1.1) \quad \gamma(\beta) = \max \left\{ 0, 1 + \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \right\}.$$

The infimum in (1.1) can be computed easily with the help of Proposition 3.2 below. In the case $\frac{a}{b} \in \mathbb{Z}$, Proposition 4.1 provides an even faster algorithm, and we are able to prove a necessary and sufficient condition for $\gamma(\beta) = 1$:

Theorem 3. *Let β be a quadratic Pisot number, root of $\beta^2 = a\beta + b$ with $a \geq b \geq 1$ and such that b divides a .*

- (i) *We have that $\gamma(\beta) = 1$ if and only if $a \geq b^2$ or $(a, b) \in \{(24, 6), (30, 6)\}$.*
- (ii) *If $a = b \geq 3$ then $\gamma(\beta) = 0$.*

This paper is organized as follows: In the next section, notions on words, representation spaces and β -tiles are recalled, and properties of β -adic expansions are studied. Section 3 connects tiles arising from the β -transformation and the value $\gamma(\beta)$ in order to prove Theorem 1. The proof of Theorem 2 is completed in Section 4, together with that of Theorem 3. Comments on the general case are in Section 5, along with a list of related open questions.

2. PRELIMINARIES

2.1. Words over a finite alphabet. We consider both finite and infinite words over a finite alphabet $\mathcal{A}$. The set of finite words over $\mathcal{A}$ is denoted $\mathcal{A}^*$. The set of all (right) infinite words over $\mathcal{A}$ is denoted $\mathcal{A}^\omega$, and it is equipped with the Cantor topology. An infinite word is (*eventually*) *periodic* if it is of the form $vu^\omega := vuuu \dots$; a finite word v is the pre-period and a non-empty finite word u is the period; if the pre-period is empty, we speak about a *purely periodic word*. A prefix of a (finite or infinite) word w is any finite word v such that w can be written as $w = vu$ for some word u . We denote by $\mathbf{u}[n]$ the prefix of length n of an infinite word $\mathbf{u}$.

To a finite word $w = w_0w_1 \dots w_{k-1}$ we assign the polynomial

$$P_w(X) := \sum_{i=0}^{k-1} w_i X^i.$$

Similarly, $P_{\mathbf{u}}(X) := \sum_{i \geq 0} u_i X^i$ is a power series for an infinite word $\mathbf{u} = u_0u_1u_2 \dots$.

2.2. Representation spaces. The following notation will be used: For integers $a, b \in \mathbb{Z}$, we denote by $a \perp b$ the fact that a and b are co-prime, i.e., that $\gcd(a, b) = 1$. Moreover, for $b \geq 2$ we put $\mathbb{Z}_b := \{p/q : p, q \in \mathbb{Z}, q \perp b\}$ (the ring of rational numbers with denominator co-prime to b).

We adopt the notation of [MS14], however, we restrict ourselves to β being a quadratic Pisot number. Let $K = \mathbb{Q}(\beta)$. Since β is quadratic, there are exactly two infinite places of K ; they are given by the two Galois isomorphisms of $\mathbb{Q}(\beta)$: the identity and $x \mapsto x'$ that maps β to its Galois conjugate. Both these places have $\mathbb{R}$ as their completion.

If β is not a unit, then we have to consider finite places of K as well. We define the ring K_f as the direct product $K_f := \prod_{\mathfrak{p} | (\beta)} K_{\mathfrak{p}}$, where $\mathfrak{p}$ runs through all prime ideals of $\mathbb{Q}(\beta)$ that divide the principal ideal (β) and $K_{\mathfrak{p}}$ is the associate completion of $\mathbb{K}$; for a precise definition, we refer to [MS14, §2.2]. The direct products $\mathbb{K} := K \times K' \times K_f$ and $\mathbb{K}' := K' \times K_f$ are called *representation spaces*. We define the diagonal embeddings

$$\delta: \mathbb{Q}(\beta) \rightarrow \mathbb{K}, x \mapsto (x, x', x_f) \quad \text{and} \quad \delta': \mathbb{Q}(\beta) \rightarrow \mathbb{K}', x \mapsto (x', x_f),$$

where x_f is the vector of the embeddings of x into the spaces $K_{\mathfrak{p}}$. We put

$$S_f := \overline{\{x_f : x \in S\}} \quad \text{for any } S \subseteq K.$$

In particular, we consider $\mathbb{Z}[\beta]_f$, which is a compact subset of K_f . Since multiplication by β_f is a contraction on K_f , we have that $\beta_f^n \mathbb{Z}[\beta]_f \rightarrow \{0_f\}$ as $n \rightarrow \infty$.

2.3. Beta-tiles. For $x \in [0, 1)$, we define the (reflected and translated) β -tile of x as the Hausdorff limit

$$\mathcal{Q}(x) := \lim_{k \rightarrow \infty} \delta'(x - \beta^k T^{-k}(x)) \subseteq \mathbb{K}'.$$

Note that the standard definition of a β -tile for $x \in \mathbb{Z}[\beta^{-1}] \cap [0, 1)$ is $\mathcal{R}(x) := \delta'(x) - \mathcal{Q}(x)$, see e.g. [MS14]. For a quadratic Pisot number β , root of $\beta^2 = a\beta + b$ with $a \geq b \geq 1$, we have that $\mathcal{Q}(x) = \mathcal{Q}(0)$ for $x < \beta - a$ and $\mathcal{Q}(x) = \mathcal{Q}(\beta - a)$ otherwise. The dynamical system $([0, 1), T)$ admits $(\mathcal{X}, \mathcal{T})$ as its natural extension, where

$$\mathcal{X} := ([0, \beta - a) \times \mathcal{Q}(0)) \cup ([\beta - a, 1) \times \mathcal{Q}(\beta - a)) \subset \mathbb{K}$$

is a union of two suspensions of β -tiles and $\mathcal{T}(x, y) := \delta(\beta)(x, y) - \delta(\lfloor \beta x \rfloor)$. The natural extension domain is often required to be a closed set, but here it is more convenient to work with the one above, since the following result holds:

Theorem B ([HI97, IR05, BS07]). *For a Pisot number β , we have that x has a purely periodic β -expansion if and only if $x \in \mathbb{Q}(\beta)$ and $\delta(x) \in \mathcal{X}$.*

2.4. Beta-adic expansions. In Definition 1.2, β -adic expansions are defined on $\mathbb{Z}[\beta]$. By Lemma 2.2 below, we extend this definition to the closure $\mathbb{Z}[\beta]_f$ similarly to the p -adic case. To this end, let

$$D: \mathbb{Z}[\beta]_f \rightarrow \mathbb{Z}[\beta]_f, \quad x \mapsto \beta_f^{-1}(x - d(x)_f),$$

where $d(x)$ is the unique digit $d \in \mathcal{A} := \{0, 1, \dots, |N(\beta)| - 1\}$ such that $\beta_f^{-1}(x - d_f)$ is in $\mathbb{Z}[\beta]_f$. Such d exists because $\mathbb{Z}[\beta] = \mathcal{A} + \beta\mathbb{Z}[\beta]$. It is unique because $(c + \beta\mathbb{Z}[\beta])_f \cap (d + \beta\mathbb{Z}[\beta])_f \neq \emptyset$ implies $(\beta^{-1}(c - d))_f \in \mathbb{Z}[\beta]_f$ and thus $c \equiv d \pmod{N(\beta)}$ by the following lemma:

Lemma 2.1 ([MS14, Lemma 5.2 and Eq. (5.1)]). *For each $x \in \mathbb{Z}[\beta^{-1}] \setminus \mathbb{Z}[\beta]$ we have $x_f \notin \mathbb{Z}[\beta]_f$. There exists $k \in \mathbb{N}$ such that $\mathbb{Z}[\beta^{-1}] \cap \beta^k \mathcal{O} \subseteq \mathbb{Z}[\beta]$, where $\mathcal{O}$ is the ring of integers in $\mathbb{Q}(\beta)$.*

Lemma 2.2. *The β -adic expansion map $\mathbf{h}_f: \mathbb{Z}[\beta]_f \rightarrow \mathcal{A}^\omega$ defined by*

$$\mathbf{h}_f(z) := u_0 u_1 u_2 \cdots, \quad \text{where } u_i := d(D^i(z)),$$

is a homeomorphism. It satisfies that $\mathbf{h}_f(x_f) = \mathbf{h}(x)$ for all $x \in \mathbb{Z}[\beta]$.

Proof. The map $\mathbf{h}_f$ is surjective because $\mathbf{h}_f(P_{\mathbf{u}}(\beta_f)) = \mathbf{u}$ for all $\mathbf{u} \in \mathcal{A}^\omega$. It is injective because $\mathbf{h}_f(z) = \mathbf{u} = u_0 u_1 u_2 \cdots$ implies that $z \in \sum_{i=0}^{n-1} u_i \beta_f^i + \beta_f^n \mathbb{Z}[\beta]_f$ for all n , thus $z = P_{\mathbf{u}}(\beta_f)$.

Since $\mathcal{O}_f$ is open and $\mathbb{Z}[\beta^{-1}]_f = K_f$, we get from Lemma 2.1 that $\mathbb{Z}[\beta]_f = \bigcup_{x \in \mathbb{Z}[\beta]} x_f + \beta_f^k \mathcal{O}_f$ for some $k \in \mathbb{N}$, and therefore it is an open set as well. Then the preimage $\mathbf{h}_f^{-1}(v\mathcal{A}^\omega) = P_v(\beta_f) + \beta_f^n \mathbb{Z}[\beta]_f$ is open for any $v \in \mathcal{A}^*$. As the cylinders $\{v\mathcal{A}^\omega : v \in \mathcal{A}^*\}$ form a base of the topology of $\mathcal{A}^\omega$, the map $\mathbf{h}_f$ is continuous.

The inverse $\mathbf{h}_f^{-1}$ is continuous because $\beta_f^n \mathbb{Z}[\beta]_f \rightarrow \{0_f\}$ as $n \rightarrow \infty$.

For $x \in \mathbb{Z}[\beta]$, the equality $\mathbf{h}_f(x_f) = \mathbf{h}(x)$ follows from the fact that $\beta^{-1}(x - d(x_f)) \in \mathbb{Z}[\beta]$. $\square$

Note that we can also identify $\mathbb{Z}[\beta]_f$ with the inverse limit space $\varprojlim \mathbb{Z}[\beta]/\beta^n \mathbb{Z}[\beta]$. Indeed defining the isomorphism

$$\kappa: \mathcal{A}^\omega \rightarrow \varprojlim \mathbb{Z}[\beta]/\beta^n \mathbb{Z}[\beta], \quad u_0 u_1 u_2 \cdots \mapsto (\xi_1, \xi_2, \xi_3, \dots), \quad \text{where } \xi_n = \sum_{i=0}^{n-1} u_i \beta^i,$$

the following diagram commutes:

$$\begin{array}{ccc} \mathbb{Z}[\beta]_f & \xrightarrow{D} & \mathbb{Z}[\beta]_f \\ \parallel \mathbf{h} & & \parallel \mathbf{h} \\ \mathcal{A}^\omega & \xrightarrow{(\text{shift})} & \mathcal{A}^\omega \\ \parallel \kappa & & \parallel \kappa \\ \varprojlim \mathbb{Z}[\beta]/\beta^n \mathbb{Z}[\beta] & \longrightarrow & \varprojlim \mathbb{Z}[\beta]/\beta^n \mathbb{Z}[\beta] \end{array}$$

3. BETA-TILES AND THE VALUE $\gamma(\beta)$

The goal of this section is to prove Theorems 1 and 2, using the connection between β -tiles and the value of $\gamma(\beta)$. First we prove the following lemma about the closures of $\mathbb{Z}$ and $\mathbb{Z}_b$ in K_f :

Lemma 3.1. *We have that $(\mathbb{Z})_f = (\mathbb{Z}_b)_f = (\mathbb{Z}_b \cap [c, d])_f$ for all $c < d$.*

Proof. We have that $(\mathbb{Z}_b)_f = (\mathbb{Z}_b \cap [c, d])_f$ by [ABBS08, Lemma 4.7]. Clearly $\mathbb{Z} \subseteq \mathbb{Z}_b$ whence $(\mathbb{Z})_f \subseteq (\mathbb{Z}_b)_f$. We will prove that $(\mathbb{Z}_b)_f \subseteq (\mathbb{Z})_f$, namely that every point $x/q \in \mathbb{Z}_b$ for $x, q \in \mathbb{Z}$ and $q \perp b$ can be approximated by integers. For each $n \in \mathbb{N}$, there exists $q_n \in \mathbb{Z}$ such that $q_n q \equiv 1 \pmod{b^n}$. Then $\frac{x}{q} - q_n x = (1 - q_n q) \frac{x}{q} \in \frac{1}{q} b^n \mathbb{Z} \subseteq \frac{1}{q} \beta^n \mathbb{Z}[\beta]$, therefore $(q_n x)_f \rightarrow (x/q)_f$. $\square$




FIGURE 1. The tiles $\mathcal{Q}(0)$ and $\mathcal{Q}(\beta - a)$ for $\beta = 1 + \sqrt{3}$. The (red) stripes illustrate the intersection of $Y = K' \times (\mathbb{Z})_f$ with the tiles.

Proof of Theorem 1. By Definition 1.1, Theorem B and since $\delta(1) \notin \mathcal{X}$, we have that

$$\gamma(\beta) = \inf \{ x \in \mathbb{Z}_b : x \geq 0, \delta(x) \notin \mathcal{X} \}.$$

For $x \in \mathbb{Q} \cap [0, \beta - a)$, the condition $\delta(x) \in \mathcal{X}$ is equivalent to $\delta'(x) \in \mathcal{Q}(0)$; for $x \in \mathbb{Q} \cap [\beta - a, 1)$, it is equivalent to $\delta'(x) \in \mathcal{Q}(\beta - a)$.

We recall the results of [MS14, §9.3], where the shape of the tiles is described. The intersection of $\mathcal{Q}(x)$ with a line $K' \times \{z\}$ is a line segment for any $z \in \mathbb{Z}[\beta]_f$ and it is empty for all $z \in K_f \setminus \mathbb{Z}[\beta]_f$, see Figure 1. Let $\partial^-\mathcal{Q}(x)$ denote the set of the segments' left end-points, and similarly $\partial^+\mathcal{Q}(x)$ the set of the right end-points. Put

$$l_x := \sup \pi'(\partial^-\mathcal{Q}(x) \cap Y) \quad \text{and} \quad r_x := \inf \pi'(\partial^+\mathcal{Q}(x) \cap Y) \quad \text{for } x = 0, \beta - a,$$

where $Y := K' \times (\mathbb{Z}_b)_f$ and π' denotes the projection $\pi': K' \times K_f \rightarrow K'$, $(y, z) \mapsto y$. Then all numbers $p/q \in \mathbb{Z}_b$ in $[l_0, r_0] \cap [0, \beta - a)$ have a purely periodic expansion, and so do all numbers $p/q \in \mathbb{Z}_b$ in $[l_{\beta-a}, r_{\beta-a}] \cap [\beta - a, 1)$. Outside these two sets, numbers $p/q \in \mathbb{Z}_b$ that do not have a purely periodic expansion are dense, since the points $\delta'(p/q)$ are dense in Y by Lemma 3.1. Therefore, the value of $\gamma(\beta)$ depends on the relative position of the above intervals (see Figure 1) in the following way:

$$(3.1) \quad \gamma(\beta) = \begin{cases} 0 & \text{if } l_0 > 0 \text{ or } r_0 < 0, \\ r_0 & \text{if } l_0 \leq 0 \text{ and } r_0 \in [0, \beta - a), \\ \beta - a & \text{if } l_0 \leq 0, r_0 \geq \beta - a \text{ and } \beta - a \notin [l_{\beta-a}, r_{\beta-a}], \\ \min\{r_{\beta-a}, 1\} & \text{if } l_0 \leq 0, r_0 \geq \beta - a \text{ and } \beta - a \in [l_{\beta-a}, r_{\beta-a}]. \end{cases}$$

In the rest of the proof, we will show that

$$(3.2) \quad l_0 = l_{\beta-a} - 1 = -\beta + \sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta') \quad \text{and} \quad r_0 = r_{\beta-a} = 1 + \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta').$$

As $\inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \leq P_{\mathbf{h}(0)}(\beta') = 0$, we see that (3.1) implies the statement of the theorem.

We use results of [MS14, §§8.3, 9.2 and 9.3], namely Equations (8.4) and (9.2), which read:

$$z \in \mathcal{R}(x) \cap \mathcal{R}(y) \quad \text{if and only if} \quad z = \delta'(x) + P_{\mathbf{u}}(\delta'(\beta)),$$




FIGURE 2. Boundary graph for quadratic β -tiles, cf. [MS14, Fig. 6]. Each arrow in the graph represents exactly b edges.

where $\mathbf{u} = v_0 v_1 v_2 \cdots$ is an edge-labelling of a path in the boundary graph in Figure 2 that starts in the node $y - x$; and

$$\partial \mathcal{R}(x) = \left(\mathcal{R}(x) \cap \mathcal{R}(x + \beta - \lfloor x + \beta \rfloor) \right) \cup \left(\mathcal{R}(x) \cap \mathcal{R}(x - \beta - \lfloor x - \beta \rfloor) \right),$$

where the first part is the left boundary $\mathcal{R}^-(x)$ and the second part is the right boundary $\mathcal{R}^+(x)$. Therefore

$$\begin{aligned} \partial^- \mathcal{R}(0) &= \partial^+ \mathcal{R}(\beta - a) = \mathcal{R}(0) \cap \mathcal{R}(\beta - a) = \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in (\mathcal{AB})^\omega \}, \\ \partial^+ \mathcal{R}(0) &= \mathcal{R}(a + 1 - \beta) \cap \mathcal{R}(0) = \{ \delta'(a + 1 - \beta) + P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in (\mathcal{AB})^\omega \}, \\ \partial^- \mathcal{R}(\beta - a) &= \mathcal{R}(\beta - a) \cap \mathcal{R}(2\beta - \lfloor 2\beta \rfloor) = \{ \delta'(\beta - a) + P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in (\mathcal{AB})^\omega \}, \end{aligned}$$

where we put $\mathcal{B} := \{a-b+1, a-b+2, \dots, a\}$. We have that

$$\begin{aligned} \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in (\mathcal{AB})^\omega \} &= \{ P_{(b-1)a}(\delta'(\beta)) - P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in \mathcal{A}^\omega \} \\ &= -\delta'(1) - \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in \mathcal{A}^\omega \}, \end{aligned}$$

since $\mathcal{A} = b-1-\mathcal{A}$ and $\mathcal{B} = a-\mathcal{A}$. Because $\mathcal{Q}(x) = \delta'(x) - \mathcal{R}(x)$, we have $\partial^\pm \mathcal{Q}(x) = \delta'(x) - \partial^\mp \mathcal{R}(x)$. We obtain

$$\begin{aligned} \partial^- \mathcal{Q}(0) &= \delta'(\beta - a) + \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in \mathcal{A}^\omega \}, \\ \partial^- \mathcal{Q}(\beta - a) &= \delta'(\beta - a + 1) + \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in \mathcal{A}^\omega \}, \\ \partial^+ \mathcal{Q}(0) &= \partial^+ \mathcal{Q}(\beta - a) = \delta'(1) + \{ P_{\mathbf{u}}(\delta'(\beta)) : \mathbf{u} \in \mathcal{A}^\omega \}. \end{aligned}$$

We have that

$$\delta'(1) + P_{\mathbf{u}}(\delta'(\beta)) \in Y \iff 1_{\mathbf{f}} + P_{\mathbf{u}}(\beta_{\mathbf{f}}) \in \mathbb{Z}_{\mathbf{f}} \iff P_{\mathbf{u}}(\beta_{\mathbf{f}}) \in \mathbb{Z}_{\mathbf{f}} \iff \mathbf{u} \in \mathbf{h}_{\mathbf{f}}(\mathbb{Z}_{\mathbf{f}}),$$

because $\mathbf{h}_{\mathbf{f}}(P_{\mathbf{u}}(\beta_{\mathbf{f}})) = \mathbf{u}$ and $\mathbf{h}_{\mathbf{f}}$ is a homeomorphism by Lemma 2.2. Then, since the map $\mathbb{Z}_{\mathbf{f}} \rightarrow K'$, $z \mapsto P_{\mathbf{h}_{\mathbf{f}}(z)}(\beta')$ is continuous, we get that

$$\inf \pi'(\partial^+ \mathcal{Q}(x) \cap Y) = 1 + \inf_{z \in \mathbb{Z}_{\mathbf{f}}} P_{\mathbf{h}_{\mathbf{f}}(z)}(\beta') = 1 + \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta').$$

Similarly, $\delta'(\beta - a) + P_{\mathbf{u}}(\delta'(\beta)) \in Y$ if and only if $\mathbf{u} \in \mathbf{h}_{\mathbf{f}}(\mathbb{Z}_{\mathbf{f}} - \beta_{\mathbf{f}})$, therefore

$$\sup \pi'(\partial^- \mathcal{Q}(\beta - a) \cap Y) - 1 = \sup \pi'(\partial^- \mathcal{Q}(0) \cap Y) = \beta' - a + \sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta').$$

Since $\beta' - a = -\beta$, this justifies (3.2). $\square$

Proof of Theorem 2, case $a > \frac{1+\sqrt{5}}{2}b$. Since $\beta' < 0$, we have that

$$\sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta') \leq \sup_{\mathbf{u} \in \mathcal{A}^\omega} P_{\mathbf{u}}(\beta') = P_{(b-1)0}(\beta') = \frac{b-1}{1-(\beta')^2}.$$

We will show that this quantity is $< 2\beta - a - 1$. First, we derive, using $(\beta')^2 = a\beta' + b$, $\beta = a - \beta'$ and $1 - (\beta')^2 > 0$, that it is equivalent to

$$(3.3) \quad a + ab + \beta'(a^2 + a + 2b - 2) > 0.$$

We know that $\beta < a + 1$, therefore $\beta = a + \frac{b}{\beta} > \frac{a(a+1)+b}{a+1}$ and $\beta' = -\frac{b}{\beta} > -\frac{(a+1)b}{a^2+a+b}$. As well, $a^2 + a + 2b - 2 > 0$, therefore we estimate

$$a + ab + \beta'(a^2 + a + 2b - 2) > \frac{ab^2((\frac{a}{b})^2 - \frac{a}{b} - 1) + b^2((\frac{a}{b})^2 + 2\frac{a}{b} - 2) + 2b}{a^2 + a + b}.$$

When $\frac{a}{b} > \frac{1+\sqrt{5}}{2}$, all three terms in the numerator are positive. Since the denominator is also positive, we get that $\sup_{j \in \mathbb{Z}} P_{\mathbf{h}(j-\beta)}(\beta') < 2\beta - a - 1$. Theorem 1 then implies (1.1). $\square$

The proof of the case $a \perp b$ of Theorem 2 was given in [MS14, §9]. The proof of the case $a = b$ is given in the next section on page 9, because it falls under the case when b divides a .

The following proposition shows how to compute the infimum in Theorem 2 and thus the value of $\gamma(\beta)$ in a lot of (and possibly all) cases. Comments on the computation of $\gamma(\beta)$ by Theorem 1 are in Section 5. We recall that $\mathbf{u} \llbracket n \rrbracket$ denotes the prefix of $\mathbf{u}$ of length n .

Proposition 3.2. *Let $\beta^2 = a\beta + b$ with $a \geq b \geq 2$. Then for each $n \in \mathbb{N}$ we have*

$$(3.4) \quad \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \in \min_{j \in \{0, 1, \dots, b^n - 1\}} P_{\mathbf{h}(j) \llbracket n \rrbracket}(\beta') + (\beta')^n \frac{b-1}{1-(\beta')^2} [\beta', 1].$$

Lemma 3.3. *Let $x, y \in \mathbb{Z}[\beta]$ satisfy that $x - y \in b^n \mathbb{Z}[\beta]$. Then $\mathbf{h}(x) \llbracket n \rrbracket = \mathbf{h}(y) \llbracket n \rrbracket$.*

Proof. Since $b = \beta^2 - a\beta \in \beta \mathbb{Z}[\beta]$, we have that $x - y \in \beta^n \mathbb{Z}[\beta]$. Let $\mathbf{h}(x) = u_0 u_1 \dots$. Then $x - \sum_{j=0}^{n-1} u_j \beta^j \in \beta^n \mathbb{Z}[\beta]$ and therefore $y - \sum_{j=0}^{n-1} u_j \beta^j \in \beta^n \mathbb{Z}[\beta]$, which means that $u_0 \dots u_{n-1}$ is a prefix of $\mathbf{h}(y)$. $\square$

Proof of Proposition 3.2. Set $\mu_n := \min_{j \in \{0, 1, \dots, b^n - 1\}} P_{\mathbf{h}(j) \llbracket n \rrbracket}(\beta')$. The statement actually consists of two inequalities, which will be proved separately. Let $j \in \mathbb{Z}$. Since $\mathbf{h}(j) \llbracket n \rrbracket = \mathbf{h}(j \bmod b^n) \llbracket n \rrbracket$ by Lemma 3.3 and since $\beta' < 0$, we have

$$\begin{aligned} P_{\mathbf{h}(j)}(\beta') &\geq P_{\mathbf{h}(j) \llbracket n \rrbracket (0(b-1))^\omega}(\beta') \geq \mu_n + (\beta')^{n+1} \frac{b-1}{1-(\beta')^2} & \text{if } n \text{ is even,} \\ P_{\mathbf{h}(j)}(\beta') &\geq P_{\mathbf{h}(j) \llbracket n \rrbracket ((b-1)0)^\omega}(\beta') \geq \mu_n + (\beta')^n \frac{b-1}{1-(\beta')^2} & \text{if } n \text{ is odd.} \end{aligned}$$

To prove the other inequality, let $k \in \{0, \dots, b^n - 1\}$ be such that $\mu_n = P_{\mathbf{h}(k) \llbracket n \rrbracket}(\beta')$. Then

$$\begin{aligned} P_{\mathbf{h}(k)}(\beta') &\leq P_{\mathbf{h}(k) \llbracket n \rrbracket ((b-1)0)^\omega}(\beta') = \mu_n + (\beta')^n \frac{b-1}{1-(\beta')^2} & \text{if } n \text{ is even,} \\ P_{\mathbf{h}(k)}(\beta') &\leq P_{\mathbf{h}(k) \llbracket n \rrbracket (0(b-1))^\omega}(\beta') = \mu_n + (\beta')^{n+1} \frac{b-1}{1-(\beta')^2} & \text{if } n \text{ is odd;} \end{aligned}$$

this provides the upper bound on the infimum. $\square$

4. THE CASE b DIVIDES a

In this section, we aim to prove Theorem 3, which deals with the particular case when b divides a . Table 1 shows whether $\gamma(\beta)$ is 0, 1 or strictly in between, for $b \leq 12$ and $a/b \leq 15$. The first non-trivial values are listed in Table 2. The algorithm for obtaining these values is deduced from Theorem 2 (which covers all the cases when $\frac{a}{b} \in \mathbb{Z}$ since then either $a = b$ or $a \geq 2b > \frac{1+\sqrt{5}}{2}b$), and the following proposition, which improves the statement of Proposition 3.2.

Proposition 4.1. *Let $\beta^2 = a\beta + b$ with $a \geq b \geq 2$ and $\frac{a}{b} \in \mathbb{Z}$. Then for each $n \in \mathbb{N}$ we have*

$$\inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \in \min_{j \in \{0, 1, \dots, b^n - 1\}} P_{\mathbf{h}(j) \llbracket 2n \rrbracket}(\beta') + (\beta')^{2n} \frac{b-1}{1-(\beta')^2} [\beta', 0].$$

Lemma 4.2. *Let $\beta^2 = cb\beta + b$. Let $x, y \in \mathbb{Z}[\beta]$ satisfy that $x - y \in b^n \mathbb{Z}[\beta]$ for some $n \in \mathbb{N}$. Then $\mathbf{h}(x) \llbracket 2n \rrbracket = \mathbf{h}(y) \llbracket 2n \rrbracket$. Moreover, for all $x \in \mathbb{Z}[\beta]$ and $d \in \mathcal{A}$ there exists $y \in x + b^n \mathcal{A}$ such that $\mathbf{h}(y) \llbracket 2n+1 \rrbracket = \mathbf{h}(x) \llbracket 2n \rrbracket d$.*

$a/b =$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$b = 1$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2	*	1	1	1	1	1	1	1	1	1	1	1	1	1	1
3	0	*	1	1	1	1	1	1	1	1	1	1	1	1	1
4	0	*	*	1	1	1	1	1	1	1	1	1	1	1	1
5	0	*	*	*	1	1	1	1	1	1	1	1	1	1	1
6	0	*	*	1	1	1	1	1	1	1	1	1	1	1	1
7	0	*	*	*	*	*	1	1	1	1	1	1	1	1	1
8	0	*	*	*	*	*	*	1	1	1	1	1	1	1	1
9	0	*	*	*	*	*	*	*	1	1	1	1	1	1	1
10	0	*	*	*	*	*	*	*	*	1	1	1	1	1	1
11	0	0	*	*	*	*	*	*	*	*	1	1	1	1	1
12	0	0	*	*	*	*	*	*	*	*	*	1	1	1	1

TABLE 1. The values of $\gamma(\beta)$ for the case when b divides a . The star ‘ $\star$ ’ means that the value is strictly between 0 and 1.

Proof. We have $\beta^2 = b(c\beta + 1) \in b\mathbb{Z}[\beta]$ and $b = \beta^2 - c(1 + c^2b)\beta^3 + c^2\beta^4 \in \beta^2 + \beta^3\mathbb{Z}[\beta] \subseteq \beta^2\mathbb{Z}[\beta]$, whence $\beta^2\mathbb{Z}[\beta] = b\mathbb{Z}[\beta]$ and $\beta^{2n}\mathbb{Z}[\beta] = b^n\mathbb{Z}[\beta]$ for all $n \in \mathbb{N}$. Following the lines of the proof of Lemma 3.3, we obtain that if $x - y \in b^n\mathbb{Z}[\beta]$ then $\mathbf{h}(x)$ and $\mathbf{h}(y)$ have a common prefix of length at least $2n$.

Put $\mathbf{h}(x) = u_0u_1 \dots$. Since $b^n \in \beta^{2n} + \beta^{2n+1}\mathbb{Z}[\beta]$, we have that $u_0u_1 \dots u_{2n-1}d$ is a prefix of $\mathbf{h}(x + eb^n)$ for any $e \equiv d - u_{2n} \pmod{b}$. $\square$

Proof of Proposition 4.1. We follow the lines of the proof of Proposition 3.2 for the case n even. The lower bound is the same in both statements, therefore we only need to prove that $\inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \leq P_{\mathbf{h}(k)}(\beta')$, where $k := \arg \min_{j \in \{0, \dots, b^n - 1\}} P_{\mathbf{h}(j)}(\beta')$. For each $m \in \mathbb{N}$, there exists $k_m \in \mathbb{Z}$ such that $\mathbf{h}(k_m)[[2n+2m]] \in \mathbf{h}(k)[[2n]](0\mathcal{A})^m$ by Lemma 4.2. Then

$$\inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \leq \inf_{m \in \mathbb{N}} P_{\mathbf{h}(k_m)}(\beta') \leq \inf_{m \in \mathbb{N}} P_{\mathbf{h}(k)}([n](00)^m((b-1)0)^\omega(\beta')) = P_{\mathbf{h}(k)}([n](\beta')) . \quad \square$$

Remark 4.3. We have that

$$(4.1) \quad \mu_n := \min_{j \in \{0, 1, \dots, b^n - 1\}} P_{\mathbf{h}(j)}([2n](\beta')) = \min_{j \in J_{n-1} + b^{n-1}\mathcal{A}} P_{\mathbf{h}(j)}([2n](\beta')) ,$$

where

$$J_0 := \{0\} \quad \text{and} \quad J_n := \left\{ j \in J_{n-1} + b^{n-1}\mathcal{A} : P_{\mathbf{h}(j)}([2n](\beta')) < \mu_n + |\beta'|^{2n+1} \frac{b-1}{1-(\beta')^2} \right\} .$$

To verify (4.1), we first show that the sequence $(\mu_n)_{n \in \mathbb{N}}$ is non-increasing. Let $j \in \{0, \dots, b^n - 1\}$ be such that $\mu_n = P_{\mathbf{h}(j)}([2n](\beta'))$. Then by Lemma 4.2 there exists $d \in \mathcal{A}$ such that $\mathbf{h}(j + db^n)[[2n+1]] = \mathbf{h}(j)[[2n]]0$, whence $\mu_{n+1} \leq P_{\mathbf{h}(j+db^n)}([2n+2](\beta')) \leq \mu_n$.

Suppose now that $j \in \{0, \dots, b^n - 1\} \setminus (J_{n-1} + b^{n-1}\mathcal{A})$. Then there exists $m < n$ such that $P_{\mathbf{h}(j)}([2m](\beta')) \geq \mu_m + |\beta'|^{2m+1} \frac{b-1}{1-(\beta')^2}$, therefore $P_{\mathbf{h}(j)}([2n](\beta')) > \mu_m \geq \mu_n$.

Example 4.4. As an example, the computation of $\gamma(\beta)$ for $\beta = 1 + \sqrt{3}$, the Pisot root of $\beta^2 = 2\beta + 2$, is visualized in Figure 3. For each step of the algorithm, the value of $\gamma(\beta)$ lies in the left-most interval. Already in the 5th step we obtain that $\gamma(\beta) \in [0.900834, 0.970552]$, therefore it is strictly between 0 and 1. Note that in the 9th step we have that $\mu_9 = P_{t^{(9)}}(\beta')$ with $t^{(9)} = 001100010101010001$, and $\gamma(\beta) \in [0.91012665225, 0.91587668314]$. In the 40th step, we have that $t^{(40)} = 001100(01)^4 000100(0001)^4 (00)^2 (01)^5 (00)^3 (01)^6 (00)^2 01$ and $\gamma(\beta) \approx 0.914803044$. In the

a	b	$\gamma(\beta)$	a	b	$\gamma(\beta)$
2	2	0.914803044196...	12	6	0.736114178272...
6	3	0.992963560101...	18	6	0.993897266395...
8	4	0.933542944675...	14	7	0.584906533458...
12	4	0.999897789000...	21	7	0.944526094618...
10	5	0.834150794175...	28	7	0.997984788082...
15	5	0.995306723671...	35	7	0.999986041767...
20	5	0.999999907110...	42	7	0.9999999999971...

TABLE 2. Numerical values of $\gamma(\beta)$, where $\beta^2 = a\beta + b$, that correspond to the first couple ‘ $\star$ ’ in Table 1.

200th step, we obtain

$$\gamma(1+\sqrt{3}) \approx 0.914803044196658195047293139393794152694998618733976175733141835762361.$$

Proof of Theorem 2, case $a = b$. Take $a = b \geq 4$. Then $b = \beta^2 + (b-1)\beta^3 + (2b+1)\beta^4$, therefore $\mathbf{h}(b)[4] = 001(b-1)$. According to Proposition 4.1, we have that

$$A := \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') \leq P_{001(b-1)}(\beta') = (\beta')^2 + (b-1)(\beta')^3.$$

For $a = b \geq 5$, we use the estimate $-\beta' \in (\frac{b}{b+1}, 1)$ to obtain that $A < 1 - \frac{b^3(b-1)}{(b+1)^3} < -1$, therefore $\gamma(\beta) = 0$. For $a = b = 4$, we have $P_{001(b-1)}(\beta') \approx -1.0193$, thus $A < -1$.

When $a = b = 3$, we verify that $\mathbf{h}(21)[12] = 001200020201$ and Proposition 4.1 yields $A \leq P_{001200020201}(\beta') \approx -1.0726 < -1$, therefore $\gamma(\beta) = 0$.

When $a = b = 2$, we can follow the lines of the proof of the case $a > \frac{1+\sqrt{5}}{2}b$, because we observe that (3.3) is satisfied, namely $6 + 8\beta' \approx 0.1436 > 0$. $\square$

The proof of Theorem 3 is divided into several cases.

Proof of Theorem 3, case $a \geq b^2$. Any $j \in \mathbb{Z} \setminus \{0\}$ can be written as $j = b^n(j_0 + j_1b)$, where $n \in \mathbb{N}$, $j_0 \in \mathcal{A} \setminus \{0\}$ and $j_1 \in \mathbb{Z}$. Then $\mathbf{h}(j)[2n+1] = 0^{2n}j_0$ because $b^n \in \beta^{2n} + \beta^{2n+1}\mathbb{Z}[\beta]$, whence

$$\begin{aligned} P_{\mathbf{h}(j)}(\beta') &\geq P_{\mathbf{h}(j)[2n+1]}((b-1)0)^\omega(\beta') \geq P_{0^{2n}1((b-1)0)^\omega}(\beta') \\ &= (\beta')^{2n} \left(1 + \frac{(b-1)\beta'}{1 - (\beta')^2} \right) = (\beta')^{2n} \left(1 - \frac{(b-1)b\beta}{\beta^2 - b^2} \right) > 0, \end{aligned}$$

where the last inequality was already proved in [MS14, Theorem 6]. As $\mathbf{h}(0) = 0^\omega$, we have $P_{\mathbf{h}(0)}(\beta') = 0$. From Theorem 2 we conclude that $\gamma(\beta) = 1 + \inf_{j \in \mathbb{Z}} P_{\mathbf{h}(j)}(\beta') = 1$. $\square$

The remaining cases of the proof of Theorem 3 make use of the following relations. Let $c := a/b \in \mathbb{Z}$. Then $\frac{b}{\beta^2} = \frac{1}{1+c\beta} \in 1 - c\beta + c^2\beta^2 - c^3\beta^3 + \beta^4\mathbb{Z}[\beta]$, and more generally,

$$(4.2) \quad \frac{b^n}{\beta^{2n}} \in 1 - nc\beta + \binom{n+1}{2}c^2\beta^2 - \binom{n+2}{3}c^3\beta^3 + \beta^4\mathbb{Z}[\beta] \quad \text{for any } n \in \mathbb{N}.$$

For $j = (j_0 + j_1b)b^n$ with $n \in \mathbb{N}$, and $j_0, j_1 \in \mathbb{Z}$ we have that $\frac{j}{\beta^{2n}} = j_0 \frac{b^n}{\beta^{2n}} + j_1 \beta^2 \frac{b^{n+1}}{\beta^{2n+2}}$, therefore

$$(4.3) \quad \frac{j}{\beta^{2n}} \in j_0 - j_0nc\beta + \left(j_0 \binom{n+1}{2}c^2 + j_1 \right) \beta^2 - \left(j_0 \binom{n+2}{3}c^3 + j_1(n+1)c \right) \beta^3 + \beta^4\mathbb{Z}[\beta].$$




FIGURE 3. The computation of $\gamma(1 + \sqrt{3})$. By a thick line with a bold label we denote the intervals that we ‘keep’ (these arise from numbers in J_n), by a thin line the ones that we ‘forget’. The labels next to the intervals are the corresponding prefixes $\mathbf{h}(j)[[2n]]$.

Proof of Theorem 3, case $\beta^2 = 30\beta + 6$. We have $b = 6$ and $c = 5$. As in the proof of the previous case, we will show that $P_{\mathbf{h}(j)}(\beta') \geq 0$ for all $j \in \mathbb{Z}$. Let $j \neq 0$ be written as $j = b^n(j_0 + j_1b)$ with $j_0 \in \mathcal{A} \setminus \{0\}$ and $j_1 \in \mathbb{Z}$, then $\mathbf{h}(j) = 0^{2n}u_0u_1u_2 \cdots$ for some $u_0u_1 \cdots \in \mathcal{A}^\omega$ with $u_0 = j_0$, and $P_{\mathbf{h}(j)}(\beta') = (\beta')^{2n}P_{u_0u_1 \cdots}(\beta')$. We consider the following cases:

- If $u_0 \geq 2$, then $P_{u_0 u_1 \dots}(\beta') \geq P_{2(50)^\omega}(\beta') > 0$.
- If $u_0 = 1$ and $u_1 \leq 4$, then $P_{u_0 u_1 \dots}(\beta') \geq P_{14(05)^\omega}(\beta') > 0$.
- If $u_0 u_1 = 15$, then (4.3) yields that $j_0 = 1$ and $-j_0 n c \equiv 5 \pmod{6}$, therefore $n \equiv -1 \pmod{6}$ and $n = 6n_1 - 1$, i.e., $-j_0 n c \beta = 5\beta - 30n_1\beta \in 5\beta - 5n_1\beta^3 + \beta^4\mathbb{Z}[\beta]$. Therefore

$$\frac{j}{\beta^{2n}} \in 1 + 5\beta + \left(\binom{6n_1}{2} 5^2 + j_1 \right) \beta^2 - \left(\frac{(6n_1+1)6n_1(6n_1-1)}{6} 5^3 + 30n_1j_1 + 5n_1 \right) \beta^3 + \beta^4 \mathbb{Z}[\beta].$$

The coefficient of β^3 is congruent to 0 modulo 6 regardless of the values of n_1 and j_1 .

This means that $u_3 = 0$. Then $P_{15u_20(05)^\omega}(\beta') \geq P_{1500(05)^\omega}(\beta') > 0$.

Therefore we have $P_{\mathbf{h}(j)}(\beta') \geq 0$ for all $j \in \mathbb{Z}$.

Proof of Theorem 3, case $\beta^2 = 24\beta + 6$. We have $b = 6$ and $c = 4$. We use the same technique as in the case $\beta^2 = 30\beta + 6$.

- If $u_0 \geq 2$, then $P_{u_0 u_1 \dots}(\beta') \geq P_{2(50)^\omega}(\beta') > 0$.
- If $u_0 = 1$ and $u_1 \leq 3$, then $P_{u_0 u_1 \dots}(\beta') \geq P_{13(05)^\omega}(\beta') > 0$.
- Since c is even, we get that $u_1 \equiv -j_0 n c \pmod{6}$ is even, therefore $u_0 u_1 \neq 15$.
- If $u_0 u_1 = 14$, then (4.3) gives $j_0 = 1$ and $-j_0 n c \equiv 4 \pmod{6}$, i.e., $n \equiv -1 \pmod{3}$ and $n = 3n_1 - 1$, whence $-j_0 n c \beta = 4\beta - 12n_1\beta \in 4\beta - 2n_1\beta^3 + \beta^4\mathbb{Z}[\beta]$. We derive that

$$\frac{j}{\beta^{2n}} \in 1 + 4\beta + (\text{some integer})\beta^2 - (144n_1^3 - 30n_1 + 12n_1j_1)\beta^3 + \beta^4\mathbb{Z}[\beta].$$

As above, we get that $u_3 = 0$ regardless of the values of n_1 and j_1 , thus $P_{u_0 u_1 \dots}(\beta') \geq P_{1400(05)^\omega}(\beta') > 0$. $\square$

Proof of Theorem 3, case $c := a/b < b$ and $c \notin \{4, 5\}$ when $b = 6$. Let $n := \lceil \frac{c}{b-c} \rceil$. From (4.2), the β -adic expansion $\mathbf{h}(b^n)$ starts with $0^{2n}1(nb-nc)$. If $\frac{c}{b-c} \notin \mathbb{Z}$, then we have $nb - nc > c$ and

thus $P_{1(nc)}(\beta') \leq 1 + (c+1)\beta' < 0$, using that $\beta' = -\frac{b}{\beta} < -\frac{b}{cb+1} \leq -\frac{1}{c+1}$. By Proposition 4.1, this proves that $\gamma(\beta) < 1$ if c is not a multiple of $b-c$.

Assume now that $\frac{c}{b-c} \in \mathbb{Z}$, i.e., $n = \frac{c}{b-c}$. For $j := b^n - \binom{n+1}{2}c^2b^{n+1}$, we have by (4.3) that

$$\frac{j}{\beta^{2n}} \in 1 - nc\beta - \left(\binom{n+2}{3}c^3 - \binom{n+1}{2}c^3(n+1) \right) \beta^3 + \beta^4 \mathbb{Z}[\beta].$$

Since $-nc = c - nb \in c - n\beta^2 + \beta^3 \mathbb{Z}[\beta]$ and $(n+1)c = nb \in \beta \mathbb{Z}[\beta]$, we obtain that

$$\frac{j}{\beta^{2n}} \in 1 + c\beta - \left(\binom{n+2}{3}c^3 + n \right) \beta^3 + \beta^4 \mathbb{Z}[\beta].$$

If $\binom{n+2}{3}c^3 + n \not\equiv 0 \pmod{b}$, then

$$P_{h(j)[2n+4]}(\beta') \leq P_{0^{2n}1c01}(\beta') = \frac{(\beta')^{2n+2}}{b} + (\beta')^{2n+3} = (\beta')^{2n+2} \frac{\beta - b^2}{b\beta} < 0,$$

since $1 + c\beta' = \frac{(\beta')^2}{b}$ and $\beta < a+1 \leq b^2$, therefore $\gamma(\beta) < 1$ by Proposition 4.1.

It remains to consider the case that $\binom{n+2}{3}c^3 + n \equiv 0 \pmod{b}$, i.e.,

$$n \equiv -\frac{bn(n+2)}{6}c^2n \pmod{b},$$

because $(n+1)c = nb$. Multiplying by $b-c$ gives

$$c \equiv -\frac{bn(n+2)}{6}c^3 \pmod{b}.$$

Note that $\frac{bn(n+2)}{6} = (b-c)\binom{n+2}{3} \in \mathbb{Z}$. We distinguish four cases:

- (i) If $6 \perp b$, then $c \equiv 0 \pmod{b}$, contradicting that $1 \leq c < b$.
- (ii) If $2 \mid b$ and $3 \nmid b$, then c is a multiple of $b/2$, i.e., $c = b/2$, $n = 1$. As n is also a multiple of $b/2$, we get that $b = 2$, thus $c = 1$. For $\beta^2 = 2\beta + 2$, we already know that $\gamma(\beta) < 1$, see Example 4.4.
- (iii) If $3 \mid b$ and $2 \nmid b$, then c and n are multiples of $b/3$. For $c = b/3$ we have $n \notin \mathbb{Z}$. For $c = 2b/3$, we have $n = 2$, thus $b \in \{3, 6\}$. However, $b = 6$ contradicts $2 \nmid b$ and $b = 3$ (i.e., $c = 2$) contradicts $\binom{n+2}{3}c^3 + n \equiv 0 \pmod{b}$.
- (iv) If $6 \mid b$, then c and n are multiples of $b/6$, thus $c \in \{b/2, 2b/3, 5b/6\}$, $n \in \{1, 2, 5\}$. If $n = 1$, then $b = 6$, thus $c = 3$, and $\binom{n+2}{3}c^3 + n \not\equiv 0 \pmod{b}$. If $n = 2$, then $b \in \{6, 12\}$; we have excluded that $b = 6$, $c = 4$; for $b = 12$, $c = 8$, we have $\binom{n+2}{3}c^3 + n \not\equiv 0 \pmod{b}$. If $n = 5$, then $b \in \{6, 30\}$; we have excluded that $b = 6$, $c = 5$; for $b = 30$, $c = 24$, we have $\binom{n+2}{3}c^3 + n \not\equiv 0 \pmod{b}$. $\square$

5. THE GENERAL CASE

In the general quadratic case where $1 < \gcd(a, b) < b$, the conditions of Theorem 2 need not be satisfied. This means that we have to rely on the more general Theorem 1, i.e., to compute the two values $\inf_{j \in \mathbb{Z}} P_{h(j)}(\beta')$ and $\sup_{j \in \mathbb{Z}} P_{h(j-\beta)}(\beta')$.

We can derive, in a similar manner to Proposition 3.2, that for all $n \in \mathbb{N}$,

$$(5.1) \quad \sup_{j \in \mathbb{Z}} P_{h(j-\beta)}(\beta') \in \max_{j \in \{0, 1, \dots, b^n-1\}} P_{h(j-\beta)[n]}(\beta') + (\beta')^n \frac{b-1}{1-(\beta')^2} [\beta', 1].$$

Let now $s_n \geq 1$, for $n \in \mathbb{N}$, denote the smallest positive integer such that $s_n \in \beta^n \mathbb{Z}[\beta]$, and $r_n := \frac{s_n}{s_{n-1}}$. Then $x, y \in \mathbb{Z}$ have a common prefix of length n if and only if $y - x \in s_n \mathbb{Z}$. Therefore, in both (3.4) and (5.1) we can take $\{0, 1, \dots, s_n - 1\}$ instead of $\{0, 1, \dots, b^n - 1\}$. Moreover, following Remark 4.3, we can further restrict to the sets

$$\begin{aligned} J_0 &:= \{0\}, & J_n &:= \{j \in J_{n-1} + s_{n-1}\{0, \dots, r_n - 1\} : P_{h(j)[n]}(\beta') \leq \mu_n + |\beta'|^n \frac{b-1}{1+\beta'}\}, \\ J'_0 &:= \{-\beta\}, & J'_n &:= \{j \in J_{n-1} + s_{n-1}\{0, \dots, r_n - 1\} : P_{h(j)[n]}(\beta') \geq \nu_n - |\beta'|^n \frac{b-1}{1+\beta'}\}, \end{aligned}$$

where $\mu_n := \min_{j \in \{0,1,\dots,b^n-1\}} P_{\mathbf{h}(j)[n]}(\beta')$ and $\nu_n := \max_{j \in \{0,1,\dots,b^n-1\}} P_{\mathbf{h}(j-\beta)[n]}(\beta')$.

We conclude by several open questions that arise in the study of rational numbers with purely periodic expansions:

- (A) Prove or disprove that $\gamma(\beta) = 1$ for quadratic Pisot number $\beta > 1$, a root of $\beta^2 = a\beta + b$, if and only if $\frac{a}{b} \in \mathbb{Z}$ and $a \geq b^2$ or $(a, b) \in \{(24, 6), (30, 6)\}$.
- (B) For which quadratic β we have that $\gamma(\beta) = 0$? Can we drop the restrictions on a and b in Theorem 2? More specifically, is it true that $a < \frac{1+\sqrt{5}}{2}b$ implies $\gamma(\beta) = 0$?
- (C) What is the structure of the prefixes of β -adic expansions of integers for a general quadratic β ?
- (D) What about the cubic Pisot case? Akiyama and Scheicher [AS05] showed how to compute the value $\gamma(\beta)$ for $\beta \approx 1.325$ the minimal Pisot number (or Plastic number), root of $\beta^3 = \beta + 1$. Loridant et al. [LMST13] gave the contact graph of the β -tiles for cubic units, which could be used to determine $\gamma(\beta)$ for the units, in a similar way to what Akiyama and Scheicher did. The consideration of the β -adic spaces could then allow the results to be expanded to non-units as well.

ACKNOWLEDGEMENTS

The first author acknowledges support by Grant Agency of the Czech Technical University in Prague grant SGS11/162/OHK4/3T/14 and Czech Science Foundation grant 13-03538S. The second author acknowledges support by ANR/FWF project “FAN – Fractals and Numeration” (ANR-12-IS01-0002, FWF grant I1136) and by ANR project “Dyna3S – Dynamique des algorithmes du pgcd” (ANR-13-BS02-0003).

REFERENCES

- [ABBS08] Shigeki Akiyama, Guy Barat, Valérie Berthé, and Anne Siegel, *Boundary of central tiles associated with Pisot beta-numeration and purely periodic expansions*, Monatsh. Math. **155** (2008), no. 3–4, 377–419.
- [Aki98] Shigeki Akiyama, *Pisot numbers and greedy algorithm*, Number theory (Eger, 1996), de Gruyter, Berlin, 1998, pp. 9–21.
- [AS05] Shigeki Akiyama and Klaus Scheicher, *Intersecting two-dimensional fractals with lines*, Acta Sci. Math. (Szeged) **71** (2005), no. 3–4, 555–580.
- [BS07] Valérie Berthé and Anne Siegel, *Purely periodic β -expansions in the Pisot non-unit case*, J. Number Theory **127** (2007), no. 2, 153–172.
- [HI97] M. Hama and T. Imahashi, *Periodic β -expansions for certain classes of Pisot numbers*, Comment. Math. Univ. St. Paul. **46** (1997), no. 2, 103–116.
- [IR05] Shunji Ito and Hui Rao, *Purely periodic β -expansions with Pisot unit base*, Proc. Amer. Math. Soc. **133** (2005), no. 4, 953–964.
- [IS01] Shunji Ito and Yuki Sano, *On periodic β -expansions of Pisot numbers and Rauzy fractals*, Osaka J. Math. **38** (2001), no. 2, 349–368.
- [IS02] ———, *Substitutions, atomic surfaces, and periodic beta expansions*, Analytic number theory (Beijing/Kyoto, 1999), Dev. Math., vol. 6, Kluwer Acad. Publ., Dordrecht, 2002, pp. 183–194.
- [LMST13] Benoît Loridant, Ali Messaoudi, Paul Surer, and Jörg M. Thuswaldner, *Tilings induced by a class of cubic Rauzy fractals*, Theoret. Comput. Sci. **477** (2013), 6–31.
- [MS14] Milton Minervino and Wolfgang Steiner, *Tilings for Pisot beta numeration*, Indag. Math. (N.S.) **25** (2014), no. 4, 745–773.
- [Rén57] Alfréd Rényi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar **8** (1957), 477–493.
- [Sch80] Klaus Schmidt, *On periodic expansions of Pisot numbers and Salem numbers*, Bull. London Math. Soc. **12** (1980), no. 4, 269–278.

DEPT. MATH. FNSPE, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TROJANOVA 13, PRAGUE 12000, CZECH REP.
LIAFA, CNRS UMR 7089, UNIVERSITÉ PARIS DIDEROT – PARIS 7, CASE 7014, 75205 PARIS CEDEX 13, FRANCE
E-mail address: tohecz@gmail.com

LIAFA, CNRS UMR 7089, UNIVERSITÉ PARIS DIDEROT – PARIS 7, CASE 7014, 75205 PARIS CEDEX 13, FRANCE
E-mail address: steiner@liafa.univ-paris-diderot.fr